

Valproic Acid

Valproic acid (Depakote), or valproate, is an anticonvulsant and mood-stabilizing drug that works by increasing GABA concentration and blocking Na⁺channels. It is mainly indicated for epilepsy, bipolar disorder, and migraine prophylaxis. This drug increases levels of the neurotransmitter GABA by inhibiting the enzyme GABA transaminase, which acts to break down GABA. Also, valproic acid decreases repetitive neuronal firing by blocking voltage-gated Na⁺channels, inactivating them. This medication is known for causing GI distress, as well as being hepatotoxic and sometimes leading to pancreatitis. Valproic acid is contraindicated in pregnant patients, as it is teratogenic and a folate antagonist, and can lead to neural tube defects (spina bifida).



PLAY PICMONIC

Indications

First Line Tonic-Clonic Seizure Treatment

First-place Tonic-clown Caesar

Valproic acid is a first line drug for tonic-clonic seizures and can also be used for absence, myoclonic and partial seizure treatment.

Bipolar Disorder

Bi-polar-bear

In addition to treating seizures, valproic acid can be used as another treatment option for manic episodes found in bipolar disorder.

Migraine Prophylaxis

Mind-rain with Purple-axes

Valproic acid may also be used for prophylaxis of migraine headaches.

Mechanism of Action

Increases GABA Concentration

Up-arrow GABA-goose

Valproic acid increases GABA concentration by inhibiting GABA transaminase, an enzyme which acts to break down GABA.

Blocks Voltage-Gated Na+ Channels

Block-guy blocking Electric-gate Salt-shaker Channel

This drug also works by suppressing repetitive neuronal activity through increased Na⁺channel inactivation; blocking and preventing Na⁺influx through fast channels.

Side Effect

GI Distress

GI with Flare-gun

Adverse effects of valproic acid include GI distress, pancreatitis, and hepatotoxicity.

Contraindication

Contraindicated in Pregnancy

Caution-tape Pregnant-woman

Valproic acid is contraindicated during pregnancy because of its teratogenic properties and association with neural tube defects in the fetus (spina bifida), as it is a folate antagonist.