

## Hypersensitivity Type IV

Type IV hypersensitivity, also known as delayed-type hypersensitivity, involves T-cell activation after being presented antigens by antigen presenting cells. Macrophages are activated and cytokines are released, leading to tissue injury.



PLAY PICMONIC

### Delayed-Type Hypersensitivity

#### Delayed-sign

Type IV hypersensitivity is also known as delayed-type hypersensitivity, which leads to immune inflammation 1-10 hours after exposure to the antigen.

### T Cell Mediated

#### Tennis Ball

This hypersensitivity is T-cell mediated, and involves  $CD4^+$  and  $CD8^+$  cells. Antigen presenting cells (APCs) cause  $CD4^+$  cells to secrete cytokines, activate phagocytes, and recruit neutrophils, which leads to tissue injury. Some types of delayed-type hypersensitivity signal the  $CD8^+$  cells to directly kill tissue.

### Antibody Independent

#### Ant-tie-body Independence-protest

It is important to know that this process is antibody independent. T lymphocytes encounter antigens by APCs and directly react.

### Macrophages Activated

#### Mac-men Activated

After encountering APCs (antigen-presenting cells),  $CD4^+$  T Cells activate macrophages, which serve to eliminate the offending antigen. The cost of this, however, is sustained, continued inflammation and tissue injury.

### Examples

#### Transplant Rejection

##### Train-plant Derailed

Delayed hypersensitivity also plays a major role in chronic transplant rejection as a result of cytotoxic T cell destruction of donor cells (host versus graft rejection) or recipient cells (graft versus host rejection).

#### PPD Skin Test

##### Pea-PD officer

The tuberculin reaction, Mantoux test, or PPD, where the protein-containing antigen of the tuberculin bacillus is injected intracutaneously. In previously exposed patients induration and reddening occurs within 8-12 hours and peaks at 24-72 hours. It occurs because of accumulation of

mononuclear cells, CD4+ T cells and macrophages.

### **Contact Dermatitis**

[Contact-sports Dermatologist examining rash](#)

Contact dermatitis is another example of DTH (delayed-type hypersensitivity), and can occur from contact with the antigenic component of poison ivy or oak. It causes vesicular dermatitis in areas of contact.