

## Acute Respiratory Distress Syndrome (ARDS) Interventions



PLAY PICMONIC

### Interventions

#### Closely Monitor Patient

##### Monitor Close to Patient

Patients with ARDS are at an increased risk for developing renal failure and stress ulcers. Close monitoring of patients for these conditions is essential. Hemodynamic monitoring should also be an important component of care, as these patients may experience hypotension, hypoxemia, and hypercapnia, which can cause negative consequences if left untreated.

#### ABG's (Arterial Blood Gases)

##### AirBaG

Arterial blood gases should be checked frequently to monitor the patient's gas exchange and his/her response to oxygen therapy. In patients with ARDS, PaO<sub>2</sub> should be kept at or above 60 mmHg.

#### Oxygen

##### O<sub>2</sub>-tank

The goal of oxygen therapy is to correct hypoxemia and achieve a PaO<sub>2</sub> of 60 mmHg or higher. Patients should be treated with the lowest concentration of oxygen possible to stay within the desired parameters.

#### Assess for O<sub>2</sub> Toxicity

##### Assess-man O<sub>2</sub>-tank in Toxic-green-glow

Administration of oxygen with an FiO<sub>2</sub> greater than 60 percent for 2 days or more increases a patient's risk of developing oxygen toxicity. Signs and symptoms of oxygen toxicity include change in vision, ringing in the ears, nausea, twitching, and irritability.

#### Mechanical Ventilation

##### Machine Vent ventilating

Patients with ARDS need to be mechanically ventilated in order to maintain adequate levels of oxygenation. Long-term ventilation increases a patient's risk of developing pneumonia. Frequent mouth care and rigid infection control measures should be implemented to decrease the risk of this complication.

#### PEEP

##### Peep-marshmallow

Positive end expiratory pressure (PEEP) is applied, using a ventilator, to prevent alveolar collapse upon expiration.

## Assess for Pneumothorax

### Assess-man and Nude-Thor-axe

Due to decreased lung compliance and high peak airway pressures, barotrauma or trauma to the lung(s) is common in ventilated patients with ARDS. To reduce the risk of injuring the lung, smaller tidal volumes should be used with these patients.

## Permissive Hypercapnia

### Permitted Hiker-cape

Smaller tidal volumes are recommended when ventilating these patients. This intervention causes a rise in PaCO<sub>2</sub>. Permissive hypercapnia is allowed in this situation and is usually tolerated well.

## Considerations

## High Mortality Rate

### Up-arrow Death

The mortality rate for patients with ARDS is about 50 percent; however, prone positioning has shown to increase positive outcomes in some patients. When a patient is supine, fluid will pool in the dependent areas of the lungs, damaging alveoli, and interfering with gas exchange. Placing the patient in prone position will allow previously dependent areas of the lungs to continue to receive the greatest blood flow, and alveoli will be allowed to reopen, due to the shifting of fluid to the anterior surface of the lungs.