

# Busulfan

Busulfan is an alkylating anticancer agent used to treat CML and as a conditioning agent prior to bone marrow transplantation. As an alkylating agent, it works to crosslink DNA. Side effects of busulfan use include myelosuppression, pulmonary fibrosis, and hyperpigmentation.



**PLAY PICMONIC** 

#### **Indications**

#### **CML**

CaMeL

Busulfan is used to treat CML (chronic myelogenous leukemia) and because of its low cost. The gold standard of treatment for CML, now, is imatinib.

#### **Bone Marrow Ablation**

Bone Arrow Baster

This drug is used to ablate a patient's bone marrow and is indicated as a conditioning agent prior to bone marrow transplantation, especially in patients with CML.

### Mechanism

### Cross-Links DNA

Linked DNA

Busulfan is an alkylating agent that forms DNA-DNA intrastrand crosslinks. This process prevents DNA replication, as these crosslinks cannot be repaired by cellular machinery, causing the cancer cell to undergo apoptosis.

### **Side Effects**

### Myelosuppression

Suppressed Red and White-blood-cells

This drug leads to severe myelosuppression, hence its use as an ablation agent.

## **Pulmonary Fibrosis**

Fiber-ball hitting Lungs

The most notable toxicity of this medication is interstitial pulmonary fibrosis, leading to the term "Busulfan lung."

### Hyperpigmentation

Hiker-pig with Hyperpigmentation

Patients can develop skin hyperpigmentation with the use of Busulfan. The presentation of this skin manifestation mimics Addison's disease, affecting the joints and skin creases as bronzing.