

## Alzheimer's Disease Assessment (Late Symptoms)

Dementia is a progressive deterioration of cognitive functioning and global impairment of the intellect. Alzheimer's disease (AD) is one of the most common forms of dementia characterized by neuronal degeneration and brain cell death. Late signs and symptoms of this disease process include memory impairment, wandering behavior, confabulation, perseveration, lack of abstract thinking, and severe cognitive decline.



PLAY PICMONIC

### Late Signs and Symptoms

#### Memory Impairment

##### Im-peared Memory

As Alzheimer's progresses, patients display little memory and are unable to process new information. Furthermore, they may not remember knowing a person, lose track of time, and have impaired recent memory.

#### Wandering Behavior

##### Wandering with a Wand

At this stage in the disease, the person is easily disoriented and gets lost in familiar places, which can last for hours. Patients exhibit wandering behavior.

#### Confabulation

##### Con-fabric-tongue

Confabulation is the creation of plausible stories in place of actual memories to maintain self-esteem. Alzheimer's patients can display confabulation to providers and family members when describing their actions. Typically, family members relate to the health care provider what actually occurred and clarify any confabulated story by the patient.

#### Perseveration

##### Pear-perseveration

Perseveration describes the repetition of phrases and behaviors. This is very common among patients with advanced Alzheimer's. They may continue to focus on a particular question repeatedly or continue to repeat a gesture or task. This is because they don't remember asking or doing the task.

#### Lack of Abstract Thinking

##### Cannot think of Abstract-solution

These patients no longer have an ability to think abstractly. They cannot organize or plan accordingly, and find it difficult to draw conclusions and solve problems.

#### Severe Cognitive Decline

##### Cogs in Decline

In the final stage of Alzheimer's, the patient loses the ability to respond to the environment, carry on a conversation and may lose the ability to smile (mask-like facies). Major behavioral changes also coincide with the cognitive decline - being compulsive, having repetitive behavior, needing help with

eating and using the toilet.