picmonic

Cromolyn (Mast Cell Stabilizers)

Cromolyn sodium (Intal) or cromoglicic acid is a medication that prevents the breakdown of mast cells. By preventing their breakdown, this medication prevents the release of inflammatory agents like histamine from entering the tissue. This medication is now often given second line in asthma prophylaxis, because it is required to be given 4 times per day on a fixed schedule to be effective.
br/>



PLAY PICMONIC

Mechanism

Mast Cell Stabilizer

Mast Sail Stabilized

This medication stabilizes mast cells. Mast cells break down as part of the inflammatory reaction and release histamine, a powerful vasodilator. These medications prevent mast cells from breaking down and releasing their contents.

Indications

Asthma Prophylaxis

Asthma-inhaler with Purple-axes

This medication is indicated only for the prevention of asthma attacks, as it prevents the breakdown of mast cells. It has no indication for treatment of an acute asthma attack.

Bronchial Inflammation

Broccoli in Flames

Cromolyn is used to prevent exercise-induced bronchospasm by working as an antiinflammatory. It should be administered approximately 15 minutes prior to exercise.

Allergic Rhinitis

Allergy-alligator with Runny-nose

Cromolyn is indicated as a nasal spray in some countries for the treatment of allergic rhinitis caused by irritants. It prevents the body from initiating an inflammatory reaction.

Side Effects

Cough

Coughing Coffee-pot

Patients may experience a cough as a side effect of the medication, because it increases the viscosity of mucus secretions.

Bronchospasm

Broccoli-spaceship

In very rare instances, patients have a paradoxical bronchospasm when taking this medication.

Consideration



Fixed Schedule Administration

Fixed-clock Administered

Because this medication is used to prevent inflammatory responses like asthma, it must be maintained at a constant therapeutic level. This medication must be given on a fixed schedule and is administered via nebulizer or a metered dose inhaler (MDI). Because of administration 4 times daily, other medications are now preferred for use in asthma prophylaxis.