

Hypercholesterolemia Risk Stratification

In order to properly treat patients with hypercholesterolemia, one must take into account a patient's serum LDL levels in conjunction with their risk factors and 10-year risk of having a coronary heart disease (CHD)-related event, such as a myocardial infarction. The higher the number of risk factors a patient has, the lower their target LDL level must be to prevent future adverse events. Doctors may suggest lifestyle modifications or pharmacotherapy to lower LDL levels and a patient's risk of CHD-related events.



PLAY PICMONIC

0 - 1 Risk Factors

(0) Zorro with (1) Wand Risk

In patients with 0-1 risk factors for coronary heart disease or in those with a calculated risk 10-year risk of CHD-related events $<10\%$, target LDL is <160 mg/dL.

160 mg/dL LDL goal

Less-than Sweet (16) Lard-Devil

Lifestyle modifications are recommended for those with LDL between 160 and 190 mg/dL; pharmacotherapy should be recommended in this population once LDL >190 mg/dL.

greater or equal to 2 Risk Factors and 10-year Risk $\leq 20\%$

\geq (2) Tutu Risk at (10) Tin with \leq (20) Dollar-bill

In patients with 2 or more risk factors for coronary heart disease or in those with a calculated 10-year risk of CHD-related events between 10 and 20%, target LDL is <130 mg/dL.

130 mg/dL LDL Goal

Less-than Friday the (13th) with Lard-Devil

Pharmacotherapy is recommended in this population to achieve the LDL goal of <130 mg/dL.

Coronary Heart Disease OR \geq 2 Risk Factors AND 10-Year Risk $> 20\%$

Heart Diseased OR \geq (2) Tutu Risk at (10) Tin with $>$ (20) Dollar-bill

For patients with coronary heart disease or a risk-equivalent condition (peripheral vascular disease, diabetes mellitus, aortic stenosis, etc.) or 2 or more risk factors and a calculated 10-year risk of CHD-related events $>20\%$, target LDL is <100 mg/dL.

100 mg/dL LDL Goal

Less-than (100) Dollar-bill with Lard-Devil

Pharmacotherapy is necessary in this population to achieve the LDL goal of <100 mg/dL.