

Salmonella Overview

Salmonella spp. is a gram negative bacilli. It is a non-lactose fermenter and produces white colonies on MacConkey agar. It is oxidase negative and produces hydrogen sulfide. Its flagella provides motility to facilitate invasion of Peyer's patches within the ileum. Some of the clinical conditions it causes include gastroenteritis, osteomyelitis in sickle cell patients, and reactive arthritis. Antibiotics are not routinely recommended for treatment of mild disease.



PLAY PICMONIC

Pathophysiology

Gram-Negative Rod

[Graham-cracker Negative-devil with Rod](#)

These bacilli bacteria lack the thick peptidoglycan layer of gram positives that normally retain crystal violet dye; when dyed with a Gram counterstain their thin peptidoglycan layer appears pink on microscopy.

Oxidase Negative

[Wilting Ox-daisy](#)

An oxidase test is used to determine if bacteria produce certain cytochrome c oxidases to help differentiate similar bacteria.

Non Lactose Fermenter

[Nun with Milk Fern](#)

MacConkey agar is a lactose-containing selective medium which inhibits the growth of gram positive bacteria and promotes gram negative bacteria growth. Salmonella cannot ferment the lactose in the medium, so the colonies that grow appear white.

White on Macconkey Agar

[White Monkey Petri-dish](#)

When non-lactose fermenting bacteria grow on MacConkey agar, they must use peptones as their energy source; when processing peptones, they form basic ammonia, raising the pH of the agar, and forming white colonies.

H2S Producing

[H-shaped Sulfur-match with H2S-fart](#)

These bacteria can oxidize organic compounds for energy while reducing sulfate (SO_4^-) to hydrogen sulfide (H_2S) as a paired redox reaction. This characteristic helps distinguish Salmonella from Shigella, which is non-hydrogen sulfide producing.

Flagella

[Flag](#)

Flagella are long tails made of proteins like flagellin with an internal proton pump motor that facilitates bacterial motility.

Invades GI Tract

[GI Tract](#)

Salmonella bacteria spread via the fecal-oral route; once ingested they colonize the small intestine and invade the body through Peyer's patches located in the terminal ileum. A large amount of Salmonella bacteria must be ingested because stomach acid normally destroys the bacteria.

Peyer's Patches

[Pirate Patch](#)

Peyer's patches are aggregates of lymphoid tissue in the terminal ileum of the small intestine. Peyer's patches are lined by M cells, which are specialized epithelial cells that endocytose antigens and then present them to other immune cells like T-cell lymphocytes.

Considerations

Osteomyelitis in Sickle Cell Patients

[Skeleton-on-fire with Sickle](#)

Due to its encapsulated trait that requires a functional spleen to properly destroy, Salmonella infection is classically associated with osteomyelitis in patients with sickle cell anemia who often have nonfunctional spleens.

Reactive Arthritis

[Reaction King-Arthur](#)

Some patients develop reactive arthritis after a Salmonella infection. Reactive arthritis consists of the classic triad: uveitis, urethritis, and arthritis. Its clinical manifestations can be remembered by the saying "Can't see, can't pee, can't climb a tree."

Antibiotics Prolong Fecal Excretion

[ABX-guy causing Prolonged Feces](#)

The use of antibiotics to treat salmonellosis is debated, as antibiotic treatment has been associated with a prolonged carrier state and symptoms. Supportive care is recommended and most infections are self-limiting without antimicrobial use.