

Drug Trial Phases

There are five sequential phases in a drug trial, each with a different purpose and population size. The phases start from small scale Phase 0 trials which discern pharmacological properties in healthy volunteers, followed by Phase I studies where safety and response in patients with the disease of interest is evaluated. In Phase II, efficacy and side effect identification is completed, leading to Phase III trials where the new drug is compared with standard of care or placebo through randomized control trials. Finally, Phase IV trials represent post-marketing studies conducted after a drug is approved for use by the FDA for surveillance and evaluation of pharmacoeconomics.



PLAY PICMONIC

Phase 0

Phase 0

[Phase \(0\)](#)

Phase 0 is useful to eliminate drugs that don't measure up to even the most rudimentary requirements, saving time and money. Likewise it smooths the approval process to move on to the next drug trial phases in order to register a new compound.

Pharmacological Properties

[Pharmacological Property Registration](#)

The objective of Phase 0 trials is to show whether a drug's pharmacokinetics and pharmacodynamics warrant continued exploration. However, it will not demonstrate any therapeutic effect nor will it replace the dose escalation, safety, and tolerance studies required in Phase I.

Healthy Volunteers

[Healthy Volunteer](#)

In Phase 0 the drug is only given to a very small group of healthy volunteers, usually less than 15 individuals, who are administered a microdose, less than 1% of the therapeutic dose of the drug in question, for no more than seven days.

Phase I

Phase I

[Phase \(1\) Wand](#)

In Phase I trials, researchers test an experimental drug or treatment in a small group of people, around 20 to 100 people, for the first time. Around 70% of drugs move to the next phase.

Safety

[Ensure Safety Net](#)

The objective is to evaluate the drug's safety, determine a safe dosage range, and identify side effects.

Patients with Disease of Interest

[Diseased-guy](#)

Phase I studies are usually conducted in patients with the disease of interest. The goal here is to determine what the drug's most frequent side effects are and, often, how the drug is metabolized and excreted. The number of subjects typically ranges from 20 to 80.

Phase II

Phase II

[Phase \(2\) Tutu](#)

Phase II studies begin if Phase I studies don't reveal unacceptable toxicity. While the emphasis in Phase I is on safety, the emphasis in Phase II is on efficacy. Phase II usually lasts months to 2 years, and 33% of drugs move to the next phase.

Efficacy and Side Effects

[Efficacy-elephant vs Cider-effects](#)

Phase II aims to obtain preliminary data on whether the drug works in people who have a certain disease or condition. For controlled trials, patients receiving the drug are compared with similar patients receiving a different treatment, usually an inactive substance (placebo), or a different drug. Safety continues to be evaluated, and short-term side effects are studied. Typically, the number of subjects in Phase II studies ranges from a few dozen to about 300.

Phase III

Phase III

[Phase \(3\) Tree](#)

Phase III studies begin if evidence of efficacy is shown in Phase II. Phase III trials evaluate efficacy using more subjects and comparing the treatment group vs the standard treatment or vs placebo. The number of subjects usually ranges from several hundred to about 3,000 volunteers. Phase III usually lasts 1 to 4 years. Only 25-30% of the drugs move to phase IV.

Comparison with Standard of Care or Placebo

[Standard Care-bear and Sugar-pill](#)

The purpose of Phase III is to evaluate how the new medication works in comparison to existing medications for the same condition.

Randomized Control Trial

[Randomized Controller](#)

Comparisons are achieved through double-blind randomization of volunteers into two groups where they receive the novel drug vs the comparison existing medication. Double-blind methodology helps to eliminate bias when interpreting results.

Phase IV

Phase IV

[Phase \(4\) Fork](#)

Phase IV trials are post-marketing studies conducted after a drug is approved for use by the FDA. They are usually large studies, including several thousand volunteers with the disease/condition.

Surveillance

[Surveillance-camera](#)

The goal of Phase IV trials is to provide additional information including the treatment or drugs risks, benefits, and best use.

Pharmacoeconomics

[Pharmaceuticals + Money Bag Prize](#)

These post-marketing requirement and commitment studies are sponsored and conducted after the FDA has approved a product for marketing. The FDA uses these studies to gather additional information about a product's safety, efficacy, or optimal use.