

Tricyclic antidepressants can be harmful in elderly patients due to side effects such as sedation and orthostatic hypotension. SSRIs can be used as an antidepressant in the elderly with caution.

NSAIDs

N-sad

NSAIDs are included due to the risk of GI bleeding. They should also be avoided in symptomatic heart failure patients and high-risk groups such as those aged >75 years old or patients taking corticosteroids, anticoagulants, or antiplatelet drugs. Meloxicam is known to have more CNS side effects compared with other NSAIDs.

Antispasmodics

Ant-tie on Spaceship

Certain antispasmodics should be avoided in the elderly due to their anticholinergic effects. However, they still can be administered for short-term palliative care.

Opioids

Poppy-droid

The reduction of CYP3A4's function to clear opioids from the body can be seen in elderly patients. Because of this, the use of opioids should be cautioned. Opioids should not be used alongside benzodiazepines due to their risk of overdose. Gabapentin/pregabalin should also be avoided with opioids due to its sedation-related side effects.

Alpha Blockers

Afro Block-guy

Alpha-blockers are used for the treatment of hypertension. Orthostatic hypotension may occur in the elderly due to these medications. Examples of these drugs include Doxazosin, Prazosin, and Terazosin.

Antiparkinson Agents

Ant-tie Park-in-sun Garage

Elderly patients should avoid antiparkinson agents such as Benztropine and Trihexyphenidyl. These agents may be used in patients with Parkinson's, but not for the prevention of extrapyramidal symptoms in patients taking antipsychotics.