

Teriparatide



PLAY PICMONIC

PTH Analog

[Parachuting-thigh-droid with Harmonica On-a-log](#)

Teriparatide is a recombinant PTH analog commonly used in the treatment of osteoporosis and specifically approved for use in those with glucocorticoid-induced osteoporosis.

Increased Osteoblast Activity

[Up-arrow Ostrich-blast](#)

Teriparatide works to increase osteoblastic activity, which leads to bone growth. This effect occurs when administered in a pulsatile fashion.

Teriparatide has a short duration of action and half-life, and, as such, the initial effect is bone resorption. This effect is then followed by an increase in bone formation. Thus, when administered in a pulsatile fashion, this stimulates bone formation compared to resorption and, over time, leads to increased bone density.

Indications

Osteoporosis

[Ostrich-with-porous-bones](#)

As a PTH analog, teriparatide is commonly used to treat osteoporosis. It is also approved for the treatment of glucocorticoid-induced osteoporosis and can be considered a first line in higher-risk patients.

Adverse Effects

Increased Risk of Osteosarcoma

[Up-arrow Risk Ostrich-shark-comb](#)

Teriparatide should be used with caution in those with elevated risk factors for osteosarcoma. For instance, those with increased unexplained ALK-P, Paget disease of the bone, or those with a history of cancer or radiation therapy.

Hypercalcemia

[Hiker-calcified-cow](#)

Teriparatide use can result in transitory hypercalcemia.

Contraindications

Paget Disease of Bone

[Paget-disease of Bone with Caution-tape](#)

Teriparatide should be used with caution in those with elevated risk factors for osteosarcoma. For instance, those with increased unexplained ALK-P, Paget disease of the bone, or those with a history of cancer or radiation therapy.

Radiation Therapy

[Radiation-therapy with Caution-tape](#)

Teriparatide should be used with caution in those with elevated risk factors for osteosarcoma. For instance, those with increased unexplained ALK-P, Paget disease of the bone, or those with a history of cancer or radiation therapy.