

Stimulant Laxatives



PLAY PICMONIC

Mechanism of Action

Enteric Nerve Stimulation

[Intestine Nerve Stim-mule](#)

Stimulant laxatives work by triggering enteric nerve stimulation and causing colonic contraction. It takes around 6-12 hours to work.

Colonic Contraction

[Colon Flexing](#)

Stimulant laxatives will stimulate the myenteric plexus and Auerbach plexus. This stimulation will cause an increase in intestinal motility and secretions.

Indication

Constipation

[Corked Con-toilet](#)

Healthcare providers should consider pharmacological options to treat constipation after initial interventions for lifestyle and increased fiber and fluid consumption fail to relieve symptoms. The first pharmacological recommendation is bulk-forming laxatives, while osmotic laxatives and stimulant laxatives are the second lines in this regard.

Drugs

Senna

[Senna](#)

Senna is used as a short-term basis for treating constipation and can be used to empty the bowels before medical procedures, such as surgery and colonoscopy. Some brands of Senna are Senokot, Senexon, SennaGen, and Little Tummy's Stimulant Laxative Drops.

Sodium Picosulfate

[Salt-shaker Pickle-sulfur-match](#)

Patients taking sodium picosulfate are instructed to drink large amounts of water to compensate for dehydration and electrolyte imbalance that can appear. Its combination with magnesium citrate is often used to empty a patient's bowel before a colonoscopy. Some of the brand names are Laxoberon, Picolax, and Guttalax.

Bisacodyl

[Bicycle-cod](#)

Bisacodyl is also used to empty the bowels of patients before surgery due to its short-term basis. Some brands of bisacodyl are Correctol, Bisacolax, Codulax, and Dulcolax. The patient is instructed not to drink with milk and suggested to take it at bedtime.

Cascara

[Cash-mascara](#)

Cascara is one of the OTC laxatives that is now used in supplements. It can cause liver injury if used in high doses for a long period of time.

Side Effects

Diarrhea

Toilet

The most common side effect of stimulant laxatives is diarrhea. Unless indicated, it's suggested not to consume a stimulant laxative for more than a week. Other side effects may include abdominal cramps, burping, and nausea.