

## Linezolid (Zyvox)

This newer, bacteriostatic antimicrobial agent is considered part of the oxazolidinone class. Through inhibition of the 50S subunit, Linezolid is able to combat gram-positive causes of pneumonia and skin infections, as well as MRSA and VRE. This antibiotic is comparatively expensive and sometimes hard to obtain, however its use is often limited by its severe adverse effects, including bone marrow suppression and serotonin syndrome. Linezolid's efficacious use against drug-resistant organisms make it an important antibiotic to know and remember.



PLAY PICMONIC

### Mechanism Of Action

#### Oxazolidinone Antibiotic

[Ox-soldier ABX-guy](#)

Linezolid is a drug from the oxazolidinone family. It was the first of its class of antibiotics consisting of heterocyclic, nitrogen-containing, organic, 5-member ring compounds. Linezolid was first approved for use by the FDA in 2000.

#### Binds 23S rRNA of 50S Subunit

[Binds \(23\)S Jersey with rabbi-Rhino and \(50\)S Rapper Subunit](#)

Linezolid, like many other antibiotics, has a mechanism of action based around the ribosomal subunit, particularly the 50S subunit, where it binds 23S rRNA to prevent the formation of a functional 70S subunit, which is a necessary component of translation.

#### Inhibits Protein Synthesis

[Mr. Protein in Inhibiting-chains](#)

Since this drug's mechanism blocks the creation of the required 70S subunit, translational production of peptide chains cannot take place; this leads to cell stagnation and death in some cases, though Linezolid is primarily bacteriostatic.

### Indication

#### Gram-Positive Organisms

[Graham-cracker Positive-angel](#)

This medication is used to combat resistant gram-positive organisms, especially pneumonia and skin infections secondary to staphylococcus and streptococcus.

#### MRSA

[MR. Saw](#)

One of the major uses of Linezolid in today's clinical setting is as an alternative to vancomycin in fighting MRSA infections, particularly in those who may not be suitable candidates for vancomycin use.

## VRE (Vancomycin Resistant Enterococci)

### Van-tank-mice Tied-up

VRE (Vancomycin-Resistant Enterococci) are organisms that continue to be a concern due to their deleterious effects, broad drug resistance, and difficulty in being eradicated. It has been discovered that certain VRE strains are even now developing resistance to Linezolid.

## Side Effects

### Bone Marrow Suppression

#### Bone Marilyn-Monroe Suppressed

This rare, severe, adverse effect is thought to be duration-dependent (typically less than 2 weeks). Extreme caution should be used in patients who already have bone marrow suppression or chronic infections. It is recommended that the CBC is monitored weekly, and it should be discontinued in patients exhibiting evidence of myelosuppression.

### Decreased Platelet Count

#### Down-arrow Plate Count

Thrombocytopenia is the most common blood dyscrasia caused by Linezolid. Careful monitoring of platelet count and symptoms of bleeding, particularly in patients on antiplatelet therapy such as aspirin, is warranted.

### Peripheral Neuropathy

#### Purple-wavy Neuron-extremities

Sensation changes, such as numbness, tingling, burning, and itching can occur with Linezolid use. This drug is also associated with optic neuropathy, as well as diminishment or loss of proprioception, vibration, and/or temperature sensations.

### Lactic Acidosis

#### Lake of Acidic-lemon

This is one of several medications known to cause lactic acidosis in certain patients. Prescribers should be mindful of this adverse effect in patients presenting with recurrent nausea and vomiting, decreased levels of bicarbonate, or in whom an otherwise unexplained metabolic acidosis is seen.

### Serotonin Syndrome

#### Silver-tonic Savage

Exercise extreme caution for patients that exhibit a constellation of multisystem excitation symptoms indicative of serotonin syndrome. These symptoms include confusion, agitation, hallucinations, seizures, hyperthermia, shivering, tachycardia, rhabdomyolysis, and possibly organ failure in the most severe cases. Patients on Linezolid should not combine the drug with SSRIs (Selective Serotonin Reuptake Inhibitors), and they should wait 2 weeks after stopping an MAO-I (monoamine oxidase inhibitor) as serotonin syndrome can occur.

## Considerations & Patient Education

### Avoid Tyramine Containing Foods

#### Avoid Tire-man and Food

Foods such as aged cheese, fava bean pods, soy sauce, soybean-based foods, cured or aged meat, sauerkraut, beer on tap, and Marmite yeast extract should be avoided in patients taking Linezolid, as they contain high amounts of tyramine. Additionally, food at risk of being spoiled, including meat, poultry, or fish should all be avoided in patients currently taking Linezolid. Consumption of these foods can lead to tyramine excess, which may result in dangerous blood pressure changes and the associated symptoms of chest pain, tachycardia, confusion, headaches, nausea and vomiting, vision changes, epistaxis, and anxiety.